

Examples of limits of Frobenius (type) structures: the singularity case

ANTOINE DOUAI *

September 29, 2009

Abstract

We give examples of families of Frobenius type structures on the punctured plane and we study their limits at the boundary. We then discuss the existence of a limit Frobenius manifold. We also give an example of a logarithmic Frobenius manifold.

PLEASE NOTE THAT THIS PREPRINT HAS BEEN SUPERSEDED BY arXiv:0909.4063

1 Introduction

Let $w_1, \dots, w_n$ be positive integers and $f : (\mathbb{C}^*)^n \rightarrow \mathbb{C}$ be the Laurent polynomial defined by

$$f(u_1, \dots, u_n) = u_1 + \dots + u_n + \frac{1}{u_1^{w_1} \dots u_n^{w_n}}.$$

It has been explained in [7] how to attach to f a canonical Frobenius manifold: the two main ingredients are a Frobenius type structure on a point, that is a tuple

$$(E^o, R_0^o, R_\infty^o, g^o)$$

where E^o is a finite dimensional vector space over $\mathbb{C}$, g^o is a symmetric and nondegenerate bilinear form on E^o , R_0^o and R_∞^o are two endomorphisms of E^o such that $R_\infty^o + (R_\infty^o)^* = nId$ and $(R_0^o)^* = R_0^o$ (* denotes the adjoint with respect to g^o) and a pre-primitive and homogeneous section of E^o , namely a section which is a cyclic vector of R_0^o and an eigenvector of R_∞^o . The canonical solution of the Birkhoff problem for the Brieskorn lattice of f given by M. Saito's method yields the required canonical Frobenius type structure. This is the punctual construction. This gives, for $w_1 = \dots = w_n = 1$, the mirror partner of the projective space $\mathbb{P}^n$ (see [1]), and more generally the mirror partner of the weighted projective space $\mathbb{P}(1, w_1, \dots, w_n)$ (see [9] and [3]).

The purpose of these notes is to give analogous results for the deformation $F : (\mathbb{C}^*)^n \times X \rightarrow \mathbb{C}$ of f defined by

$$F(u_1, \dots, u_n, x) = u_1 + \dots + u_n + \frac{x}{u_1^{w_1} \dots u_n^{w_n}}$$

*A.Douai: Laboratoire J.A Dieudonné, UMR CNRS 6621, Université de Nice, Parc Valrose, F-06108 Nice Cedex 2, France. Email: douai@unice.fr . Partially supported by ANR grant BLAN08-1 309225 (SEDIGA)

where $X := \mathbb{C}^*$ and then to discuss the existence of a "limit" Frobenius manifold as x approaches 0. This kind of problem is also considered in [4], using another strategy (we will not use the reference [6] at all) and for a different class of functions. Notice however that the case $w_1 = \dots = w_n = 1$ is common to both papers.

Let us precise the situation: let

$$G = \frac{\Omega^n(U)[\tau, \tau^{-1}, x, x^{-1}]}{(d_u - \tau d_u F) \wedge \Omega^{n-1}(U)[\tau, \tau^{-1}, x, x^{-1}]}$$

be the (Fourier-Laplace transform of the) Gauss-Manin system of F and

$$G_0 = \frac{\Omega^n(U)[\tau^{-1}, x, x^{-1}]}{(\tau^{-1} d_u - d_u F) \wedge \Omega^{n-1}(U)[\tau^{-1}, x, x^{-1}]}$$

be (the Fourier-Laplace transform of) its Brieskorn lattice, where the notation d_u means that the differential is taken with respect to u only. G is equipped with a connection ∇ defined by

$$\nabla_{\partial_\tau}(\omega_i \tau^i) = i \omega_i \tau^{i-1} - F \omega_i \tau^i$$

and

$$\nabla_{\partial_x}(\omega_i \tau^i) = \frac{\partial \omega_i}{\partial x} \tau^i - \frac{\partial F}{\partial x} \omega_i \tau^{i+1}.$$

In particular, if we put $\theta := \tau^{-1}$, G_0 is stable under the action of $\theta^2 \nabla_{\partial_\theta}$. One defines in the same way the Gauss-Manin system (*resp.* the Brieskorn lattice) G^o (*resp.* G_0^o) of the Laurent polynomial f (see [5, section 4] for details). It turns out that one can solve the Birkhoff problem for G_0 on the whole X : G_0 is a free $\mathbb{C}[\theta, x, x^{-1}]$ -module of rank $\mu = 1 + w_1 + \dots + w_n$ and there exists a basis of G_0 in which the matrix of the connection ∇ takes the form

$$\left(\frac{A_0(x)}{\theta} + A_\infty\right) \frac{d\theta}{\theta} + \left(-\frac{A_0(x)}{\mu\theta} + R\right) \frac{dx}{x},$$

$A_0(x)$ being a $\mu \times \mu$ matrix with coefficients in $\mathbb{C}[x]$, A_∞ and R being diagonal matrices with constant coefficients (see proposition 3.1.3). This gives a Frobenius type structure on X (see [5] and [8]), that is a tuple

$$(X, E, R_0, R_\infty, \Phi, \nabla)$$

where the different objects involved satisfy some natural compatibility relations which can be extended, and this is done in section 4, to a Frobenius type structure with metric (corollary 5.1.1)

$$\mathbb{F} = (X, E, R_0, R_\infty, \Phi, \nabla, g)$$

which will be the central object of these notes. It should be emphasized that the metric g plays here a fundamental role. This Frobenius type structure, together with the data of a pre-primitive, homogeneous and ∇ -flat form, yields also a Frobenius manifold on $\Delta \times (\mathbb{C}^{\mu-1}, 0)$ where Δ denotes the open disc of radius one, centered at $x = 1$ (see [5], [8]): we will use it first to compare the canonical Frobenius manifolds attached to the different polynomials $F_x := F(\cdot, x)$, $x \in \Delta$, by the punctual construction (see section 5).

The second part of these notes (section 6) is devoted to the study of the limit, as x approaches 0, of the Frobenius type structure $\mathbb{F}$. This limit is defined using Deligne's canonical extension

$\mathcal{L}^\varphi$ such that the eigenvalues of the residue of ∇_{∂_x} are contained in $[0, 1[$: this lattice is easily described in our situation. The key point is that $gr^V(\mathcal{L}^\varphi/x\mathcal{L}^\varphi)$, the graded module associated with the Malgrange-Kashiwara V -filtration at $x = 0$, yields a Frobenius type structure on a point which can thus be seen as the canonical limit Frobenius type structure (notice that this result is not always true if we consider $\mathcal{L}^\varphi/x\mathcal{L}^\varphi$ instead of $gr^V(\mathcal{L}^\varphi/x\mathcal{L}^\varphi)$, that is if we forget the graduation). In order to define a canonical limit Frobenius manifold, we still need a pre-primitive and homogeneous section of this limit Frobenius type structure, see again [8] and the references therein: we show in section 6 that such a section exists if and only if $w_1 = \dots = w_n = 1$. In this case, we give in section 6.4 an explicit description of this canonical limit Frobenius manifold. In general, that is if there is an w_i such that $w_i \geq 2$, the situation is less clear for the following reasons: first, we do not have a general statement saying that one can derive a Frobenius manifold from the canonical limit Frobenius type structure (nevertheless, it should be emphasized that we do not assert here that such a limit does not exist); second, even if it happens to be the case, one could get several Frobenius manifolds which can be difficult to compare.

The last section is devoted to logarithmic Frobenius manifolds: if $w_1 = \dots = w_n = 1$, we show how to get, with the help now of a suitable extension of G_0 at $x = 0$, a Frobenius type structure with logarithmic pole along $\{x = 0\}$ in the sense of [10, Definition 1.6], yielding a logarithmic Frobenius manifold. If there exists a weight w_i such that $w_i \geq 2$, we have all the tools to define a Frobenius type structure with a logarithmic pole along $\{x = 0\}$, except the metric: the symmetric bilinear form constructed here is flat but not non-degenerate.

The starting point of this paper is the reference [2] in which A. Bolibruch discusses the properties of the limit of an isomonodromic family of Fuchsian systems. It happens that, in our geometric situation, this family is produced, *via* an inverse Fourier-Laplace transformation, by a solution of the Birkhoff problem for the Brieskorn lattice of a rescaling $H(u, x) = xf(u)$ of a tame regular morse function f , see [11, Chapitre VI]. This leads naturally to the following question: given a Frobenius type structure on X , what can we expect at the limit? Finally, the reference [4], and I thank C. Sabbah for a discussion about this, explains the choice of the deformation F made in these notes (the computations for the rescaling H are similar and easier to the ones performed here). The last section grew up after a discussion with C. Sevenheck who suggested me to work with the natural extensions $\mathcal{L}_0^\varphi$ of G_0 : I thank him for that.

2 The canonical solution of the Birkhoff problem for G_0^o

2.1 Preliminaries

Let $\mathcal{S}_w$ be the *disjoint* union of the sets

$$\left\{ \frac{\ell\mu}{w_i} \mid \ell = 0, \dots, w_i - 1 \right\}$$

for $i = 0, \dots, n$, where we put $w_0 = 1$. Its cardinal is equal to $\mu := 1 + w_1 + \dots + w_n$. We number the elements of $\mathcal{S}_w$ from 0 to $\mu - 1$ in an increasing way and write

$$\mathcal{S}_w = \{s_0, s_1, \dots, s_{\mu-1}\}$$

(thus, $s_k \leq s_{k+1}$). Notice that $0 \leq \frac{s_k}{\mu} < 1$. Define, for $k = 0, \dots, \mu - 1$,

$$\alpha_k = k - s_k.$$

Lemma 2.1.1 *One has $s_0 = \cdots = s_n = 0$, $s_{n+1} = \frac{\mu}{\max_i w_i}$ and $s_k + s_{\mu+n-k} = \mu$ for $k \geq n+1$.*

Proof. See [7, p. 2]. □

Corollary 2.1.2 *One has $\alpha_0 = 0, \dots, \alpha_n = n$, $\alpha_{k+1} \leq \alpha_k + 1$ for all k ,*

$$\alpha_k + \alpha_{\mu+n-k} = n$$

for $k \geq n+1$ and

$$\alpha_k + \alpha_{n-k} = n$$

for $k = 0, \dots, n$.

Proof. One has $\alpha_{k+1} \leq \alpha_k + 1$ because (s_k) is increasing. The remaining assertions are clear. □

2.2 The Birkhoff problem for G_0^o

Let A_0^o and A_∞ be the $\mu \times \mu$ matrices defined by

$$A_\infty = \text{diag}(\alpha_0, \dots, \alpha_{\mu-1})$$

and

$$A_0^o = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & \mu \\ \mu & 0 & 0 & \cdots & 0 & 0 \\ 0 & \mu & 0 & \cdots & 0 & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & \cdots & \mu & 0 \end{pmatrix}.$$

Example 2.2.1 *We will work with the following examples:*

- (1) $n = 2$ and $w_1 = w_2 = 2$: one has $\mu = 5$ and $A_\infty = \text{diag}(0, 1, 2, \frac{1}{2}, \frac{3}{2})$.
- (2) $w_1 = \cdots = w_n = 1$: one has $\mu = n+1$ and $A_\infty = \text{diag}(0, 1, \dots, n)$.

The following results are shown in [7]:

Lemma 2.2.2 (1) G_0^o is a free $\mathbb{C}[\theta]$ -module of rank μ , equipped with a connection ∇ with a pole of Poincaré rank less or equal to 1 at $\theta = 0$.

(2) The Birkhoff problem for G_0^o has a solution: there exists a basis $\omega^o = (\omega_0^o, \dots, \omega_{\mu-1}^o)$ of G_0^o over $\mathbb{C}[\theta]$ in which the matrix of the connection ∇ takes the form

$$\left(\frac{A_0^o}{\theta} + A_\infty\right) \frac{d\theta}{\theta}.$$

Moreover, the eigenvalues of A_∞ run through the spectrum at infinity of the polynomial f .

We even have a little bit more: the basis ω^o constructed in *loc. cit.* is the *canonical* solution of the Birkhoff problem given by M. Saito's method. In particular, it is compatible with the V -filtration at $\tau = 0$ (see the last assertion of [7, Proposition 3.2]). One then has $A_0^o V_\alpha \subset V_{\alpha+1}$: in other words, if $(A_0)_{ij} \neq 0$ then $\alpha_{i-1} \leq \alpha_{j-1} + 1$. One can moreover endow G_0^o with a "metric": this is discussed in section 4.

3 The Birkhoff problem for G_0

We give in this section the counterpart of the previous results for G_0 . We will put $u_0 := \frac{1}{u_1^{w_1} \dots u_n^{w_n}}$ and $\omega_0 := \frac{du_1}{u_1} \wedge \dots \wedge \frac{du_n}{u_n}$

3.1 A natural solution

Define

$$\Gamma_0 = \{(y_1, \dots, y_n) \in \mathbb{R}^n | y_1 + \dots + y_n = 1\},$$

$$\Gamma_j = \{(y_1, \dots, y_n) \in \mathbb{R}^n | y_1 + \dots + y_{j-1} + (1 - \frac{\mu}{w_j})y_j + \dots + y_n = 1\}$$

for $j = 1, \dots, n$,

$$\chi_{\Gamma_0} = u_1 \frac{\partial}{\partial u_1} + \dots + u_n \frac{\partial}{\partial u_n}$$

and, for $j = 1, \dots, n$,

$$\chi_{\Gamma_j} = u_1 \frac{\partial}{\partial u_1} + \dots + u_{j-1} \frac{\partial}{\partial u_{j-1}} + (1 - \frac{\mu}{w_j})u_j \frac{\partial}{\partial u_j} + \dots + u_n \frac{\partial}{\partial u_n}.$$

Define also, for $g = u_1^{a_1} \dots u_n^{a_n}$,

$$\phi_{\Gamma_j}(g) = a_1 \dots + a_{j-1} + (1 - \frac{\mu}{w_j})a_j + \dots + a_n$$

and

$$h_{\Gamma_j} = \chi_{\Gamma_j}(F) - F.$$

We thus have $h_{\Gamma_0} = -\mu x u_0$ and $h_{\Gamma_j} = -\frac{\mu}{w_j} u_j$ if $j = 1, \dots, n$.

Lemma 3.1.1 *One has, for any monomial g , the equality*

$$(\tau \partial_\tau + \phi_{\Gamma_j}(g))g\omega_0 = \tau h_{\Gamma_j} g \omega_0$$

in G . In particular, one has

$$\tau \partial_\tau \omega_0 = \tau h_{\Gamma_0} \omega_0.$$

Proof. Direct computation. □

This lemma is the starting point in order to solve the Birkhoff problem for G_0 , as it has been the starting point to solve the one for G_0^o (see [7, section 3]). Put $\omega_1 = u_0 \omega_0$: the equality

$$\tau \partial_\tau \omega_0 = \tau h_{\Gamma_0} \omega_0$$

becomes

$$-\frac{1}{\mu} \tau \partial_\tau \omega_0 = x \tau \omega_1.$$

Iterating the process (the idea is to define $\omega_2 = -\frac{1}{\mu}\omega_1 h_{\Gamma_1}$ etc...), one gets sections $\omega_1, \dots, \omega_{\mu-1}$ of G satisfying

$$-\frac{1}{\mu}(\tau\partial_\tau + \alpha_k)\omega_k = \tau\omega_{k+1}$$

for $k = 1, \dots, \mu - 1$ (we put $\omega_\mu = \omega_0$): this can be done as [7, section 2 and proof of proposition 3.2]. We thus have

$$\omega_k = u^{a(k)}\omega_0$$

where the multi-indices $a(k)$ are defined in [7, p. 3].

Define, for $x \in X$,

$$A_0(x) := \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & \mu \\ \mu x & 0 & 0 & \cdots & 0 & 0 \\ 0 & \mu & 0 & \cdots & 0 & 0 \\ \ddots & \cdots & \cdot & \cdots & \cdot & \cdot \\ \ddots & \cdots & \cdot & \cdots & \cdot & \cdot \\ 0 & 0 & \cdot & \cdots & \mu & 0 \end{pmatrix}$$

and

$$R = \text{diag}(0, -1, \dots, -1, -s_{\mu-1}/\mu, \dots, -s_{n+1}/\mu)$$

(the entry -1 is counted n times).

Example 3.1.2 (1) If $n = 2$ and $w_1 = w_2 = 2$, one has $R = -\text{diag}(0, 1, 1, \frac{1}{2}, \frac{1}{2})$.

(2) If $w_1 = \dots = w_n = 1$, one has $R = -\text{diag}(0, 1, 1, \dots, 1)$.

Proposition 3.1.3 (1) G_0 is a free $\mathbb{C}[x, x^{-1}, \theta]$ -module of rank μ and $\omega = (\omega_0, \dots, \omega_{\mu-1})$ is a basis of it.

(2) In the basis ω , the matrix of the connection ∇ takes the form

$$\left(\frac{A_0(x)}{\theta} + A_\infty\right)\frac{d\theta}{\theta} + \left(R - \frac{A_0(x)}{\mu\theta}\right)\frac{dx}{x}$$

where A_∞ is the diagonal matrix defined in section 2.

Proof. (1) One shows that G_0 is finitely generated as in [7, proposition 3.2], with the help of lemma 3.1.1. To show that it is free notice that, again by [7, proposition 3.2], a section of the kernel of the surjective map

$$(\mathbb{C}[x, x^{-1}])^\mu \rightarrow G_0 \rightarrow 0$$

is given by μ Laurent polynomials which vanishes everywhere. Let us show (2): the assertion about ∇_{∂_θ} is clear (by definition of the ω_k 's). Recall that the action of ∇_{∂_x} is defined, for $\eta \in G_0$, by

$$\nabla_{\partial_x}(\eta) = -\frac{\partial F}{\partial x}\eta\theta^{-1} + \partial_x(\eta) = -u_0\eta\theta^{-1} + \partial_x(\eta).$$

An easy computation shows that one has, for $\eta = u_0 u_1^{a_1} \cdots u_n^{a_n} \omega_0$,

$$u_0\eta = \frac{1}{\mu x}F\eta - \frac{1}{\mu x}\theta\left(\sum_{i=1}^n a_i - w_i\right)\eta.$$

Since $\theta^2 \nabla_{\partial_\theta}$ is induced by the multiplication by F , the matrix ∇_{∂_x} in the basis ω takes the form

$$-\frac{A_0(x)}{\mu x \theta} + \frac{1}{\mu x} T$$

where T is the diagonal matrix defined by (apply the process above to $\omega_0, \omega_1 = u^{a(1)}\omega_0, \dots, \omega_{\mu-1} = u^{a(\mu-1)}\omega_0$)

$$T_{kk} = \sum_{i=1}^n a(k-1)_i - w_i - \alpha_{k-1}.$$

Now, one has $\sum_{i=1}^n a(k-1)_i = k-2$ (see [7, section 2]) and $\sum_{i=1}^n w_i = \mu-1$ so that $T_{kk} = k-1 - \alpha_{k-1} - \mu = s_{k-1} - \mu$. Use now the symmetry property of the s_k 's (see lemma 2.1.1). Of course, $R = T/\mu$. $\square$

Remark 3.1.4 *It follows from the second part of the proposition that $(\alpha_0, \dots, \alpha_{\mu-1})$ is the spectrum at infinity of any function $F_x := F(\cdot, x)$, $x \in X$, see [7, Proposition 3.2].*

3.2 Towards the canonical extensions of G at $x = 0$

3.2.1 The φ -solution

Define $\omega_0^\varphi = \omega_0$ and $\omega_1^\varphi = xu_0\omega_0^\varphi = x\omega_1$ (such a choice is also natural because $h_{\Gamma_0} = -\mu xu_0$): one has

$$-\frac{1}{\mu}\tau\partial_\tau\omega_0^\varphi = \tau\omega_1^\varphi$$

and gets as above forms $\omega_1^\varphi, \dots, \omega_{\mu-1}^\varphi$ satisfying

$$-\frac{1}{\mu}(\tau\partial_\tau + \alpha_k)\omega_k^\varphi = \tau\omega_{k+1}^\varphi$$

for all $k = 0, \dots, \mu-2$ and

$$-\frac{1}{\mu}(\tau\partial_\tau + \alpha_{\mu-1})\omega_{\mu-1}^\varphi = x\tau\omega_0^\varphi.$$

One has also

$$\omega^\varphi = \omega P$$

where $P = \text{diag}(1, x, \dots, x)$. Put, for $x \in X$,

$$A_0^\varphi(x) = P^{-1}A_0(x)P$$

that is

$$A_0^\varphi(x) = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & \mu x \\ \mu & 0 & 0 & \cdots & 0 & 0 \\ 0 & \mu & 0 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdot & \cdots & \cdot & \cdot \\ \cdots & \cdots & \cdot & \cdots & \cdot & \cdot \\ 0 & 0 & \cdot & \cdots & \mu & 0 \end{pmatrix}$$

and

$$R^\varphi = \text{diag}(0, 0, \dots, 0, s_{n+1}/\mu, \dots, s_{\mu-1}/\mu)$$

(0 is counted $(n+1)$ -times). If $\mu = n+1$, one has $R^\varphi = 0$. $\overline{A}_0^\varphi$ will denote the value of $A_0^\varphi(x)$ at $x = 0$.

Proposition 3.2.1 (1) $\omega^\varphi = (\omega_0^\varphi, \dots, \omega_\mu^\varphi)$ is a basis of G_0 .

(2) In this basis, the matrix of the connection ∇ takes the form

$$\left(\frac{A_0^\varphi(x)}{\theta} + A_\infty\right)\frac{d\theta}{\theta} + \left(R^\varphi - \frac{A_0^\varphi(x)}{\mu\theta}\right)\frac{dx}{x}.$$

(3) For $\theta \neq 0$, the residue matrix of ∇_{∂_x} at $x = 0$ takes the form

$$R^\varphi - \frac{\overline{A}_0^\varphi}{\mu\theta}.$$

Its eigenvalues are contained in $[0, 1[$.

Proof. (1) and (2) follow from proposition 3.1.3, using lemma 2.1.1 and the fact that

$$R^\varphi \frac{dx}{x} = R \frac{dx}{x} + P^{-1} dP.$$

(3) follows because $0 \leq \frac{s_k}{\mu} < 1$. □

Remark 3.2.2 Using the variable $\tau := \theta^{-1}$, we find that the matrix of the connection ∇ takes the form

$$(-A_0^\varphi(x)\tau - A_\infty)\frac{d\tau}{\tau} + (-A_0^\varphi(x)\tau - A_\infty + H)\frac{dx}{\mu x}$$

where H is the diagonal matrix $\text{diag}(0, 1, \dots, \mu-1)$. This can be used, because the entries of H are integers, to show that the monodromies T and T' corresponding respectively to the loops around the divisors $\{\tau = 0\} \times X$ and $\mathbb{C} \times \{0\}$ in $\mathbb{C} \times X$ are related by the formula

$$T^{-1} = (T')^\mu$$

(see [4, Corollary 6.5 (ii)]).

3.2.2 The ψ -solution

Define now

- $Q = \text{diag}(1, x, \dots, x, 1, \dots, 1)$ (x is counted n -times),
- $A_0^\psi(x) = Q^{-1}A_0(x)Q$,
- $R^\psi = -\text{diag}(0, \dots, 0, \frac{s_{\mu-1}}{\mu}, \dots, \frac{s_{n+1}}{\mu})$.

Lemma 3.2.3 (1) Suppose that $\mu \geq n + 2$. One has $(A_0^\psi(x))_{1,\mu} = \mu$,

$$(A_0^\psi(x))_{i+1,i} = \mu$$

if $i \neq n + 1$ and

$$(A_0^\psi(x))_{n+2,n+1} = \mu x.$$

(2) Suppose that $\mu = n + 1$. One has $(A_0^\psi(x))_{1,\mu} = \mu x$ and

$$(A_0^\psi(x))_{i+1,i} = \mu$$

if $i = 1, \dots, n$.

Proof. Clear. □

In the sequel, $\overline{A}_0^\psi$ will denote the value of $A_0^\psi(x)$ at $x = 0$. Note that $\overline{A}_0^\psi$ is regular.

Proposition 3.2.4 (1) $\omega^\psi = \omega Q$ is a basis of G_0 ,

(2) In this basis, the matrix of ∇ takes the form

$$\left(\frac{A_0^\psi(x)}{\theta} + A_\infty\right)\frac{d\theta}{\theta} + \left(R^\psi - \frac{A_0^\psi(x)}{\mu\theta}\right)\frac{dx}{x}.$$

(3) If $\theta \neq 0$, the residue matrix of ∇_{∂_x} at $x = 0$ takes the form

$$R^\psi - \frac{\overline{A}_0^\psi}{\mu\theta}.$$

Its eigenvalues are contained in $] -1, 0]$.

Example 3.2.5 If $n = 2$ and $w_1 = w_2 = 2$, one has

$$A_0^\psi(x) = 5 \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & x & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}$$

and

$$\overline{A}_0^\psi = 5 \begin{pmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}.$$

Remark 3.2.6 Let $\mathcal{L}$ (resp. $\mathcal{L}^\varphi$, $\mathcal{L}^\psi$) be the $\mathbb{C}[x, \theta, \theta^{-1}]$ -submodule of G generated by ω (resp. ω^φ , ω^ψ). One has $\mathcal{L}^\varphi \subset \mathcal{L}^\psi \subset \mathcal{L}$. Notice that

$$\omega_0 = \omega_0^\varphi = \omega_0^\psi$$

and

$$R(\omega_0) = R^\varphi(\omega_0^\varphi) = R^\psi(\omega_0^\psi) = 0.$$

If $\mu = n + 1$, one has $\mathcal{L}^\varphi = \mathcal{L}^\psi$.

Remark 3.2.7 (*The flat basis*) Let Δ be the open disc in $\mathbb{C}$ of radius 1, centered at $x = 1$. The local basis $\omega^{flat} := \omega x^{-R}$ ($x \in \Delta$) of G_0^{an} is flat: the matrix of the connection ∇ takes the form

$$\left(\frac{A_0^{flat}(x)}{\theta} + A_\infty \right) \frac{d\theta}{\theta} - \frac{A_0^{flat}(x)}{\mu\theta} \frac{dx}{x}$$

where $A_0^{flat} = x^R A_0(x) x^{-R}$. Notice that the matrix $A_0^{flat}(x)$ has a limit $\overline{A}_0^{flat}$ when x approaches 0: this is due to the fact that the sequence (s_k) is increasing (this is equivalent to the fact that $\alpha_{k+1} \leq \alpha_k + 1$).

4 Duality

We define in this section a non-degenerate, symmetric and flat bilinear form on G_0 .

4.1

The lattice G_0^o is equipped with a non-degenerate bilinear form

$$S^o : G_0^o \times G_0^o \rightarrow \mathbb{C}[\theta]\theta^n,$$

∇^o -flat and satisfying, for $p(\theta) \in \mathbb{C}[\theta]$,

$$p(\theta)S^o(\bullet, \bullet) = S^o(p(\theta)\bullet, \bullet) = S^o(\bullet, p(-\theta)\bullet).$$

The basis $\omega^o = (\omega_0^o, \dots, \omega_{\mu-1}^o)$ given by lemma 2.2.2 is *adapted* to S^o : one has

$$S^o(\omega_k^o, \omega_j^o) = \delta_{k, n-j} \theta^n$$

for $k = 0, \dots, n$ and

$$S^o(\omega_k^o, \omega_j^o) = \delta_{k, \mu+n-j} \theta^n$$

for $k = n+1, \dots, \mu-1$. In particular, $A_\infty + A_\infty^* = nI$ and $(A_0^o)^* = A_0^o$, where $*$ denotes the adjoint with respect to S^o . All these results can be found in [7, Sect. 4].

We define, in the basis $\omega = (\omega_0, \dots, \omega_{\mu-1})$ given by proposition 3.1.3,

$$S(\omega_0, \omega_n) = x^{-1} \theta^n,$$

$$S(\omega_k, \omega_{n-k}) = x^{-2} \theta^n$$

for $k = 1, \dots, n-1$,

$$S(\omega_k, \omega_{\mu+n-k}) = x^{-1} \theta^n$$

for $k = n+1, \dots, \mu-1$ and

$$S(\omega_i, \omega_j) = 0$$

otherwise. Notice that S is constant in the flat basis ω^{flat} : one has $S(\omega_i^{flat}, \omega_j^{flat}) = S^o(\omega_i^o, \omega_j^o)$ for all i and for all j (this follows from the symmetry property of the s_k 's). Define now

$$S : G_0 \times G_0 \rightarrow \mathbb{C}[x, x^{-1}, \theta] \theta^n$$

by linearity, using the rules

$$p(\theta)(\bullet, \bullet) = S(p(\theta)\bullet, \bullet) = S(\bullet, p(-\theta)\bullet)$$

and

$$a(x)S(\bullet, \bullet) = S(a(x)\bullet, \bullet) = S(\bullet, a(x)\bullet)$$

for $p(\theta) \in \mathbb{C}[\theta]$ and $a(x) \in \mathbb{C}[x, x^{-1}]$.

If $A(\theta, x)$ denotes the matrix of the covariant derivative ∇_{∂_θ} , $\check{\nabla}_{\partial_\theta}$ will denote the covariant derivative whose matrix is $-A(-\theta, x)$ in the same basis. We will say that S is ∇ -flat if

$$\partial_\theta S(\varepsilon, \eta) = S(\nabla_{\partial_\theta}(\varepsilon), \eta) + S(\varepsilon, \check{\nabla}_{\partial_\theta}(\eta))$$

and

$$\partial_x S(\varepsilon, \eta) = S(\nabla_{\partial_x}(\varepsilon), \eta) + S(\varepsilon, \nabla_{\partial_x}(\eta)).$$

Keep the notations of section 3 and put $C_0 = -A_0(x)/\mu x$, $C_\infty = R/x$, $C_0^\varphi = -A_0^\varphi(x)/\mu x$ etc...

Lemma 4.1.1 (1) *One has*

$$(A_0(x))^* = A_0(x), \quad A_\infty + A_\infty^* = nI, \quad C_0^* = C_0,$$

where $*$ denotes the adjoint with respect to S . Same results for $A_0^{flat}(x)$, C^{flat} , $A_0^\varphi(x)$, C_0^φ , $A_0^\psi(x)$, C_0^ψ .

(2) *The bilinear form S is ∇ -flat.*

Proof. (1) The first equality follows from the definition of $A_0(x)$ and from the definition of S . For the second, use moreover the symmetry property of the numbers α_k (see lemma 2.1.2). The third equality is then clear and (2) follows from (1) and from the definition of S and R . $\square$

We have in particular

$$x\partial_x S(\varepsilon, \eta) = S(R(\varepsilon), \eta) + S(\varepsilon, R(\eta))$$

and this give a symmetry property for the matrix R . Notice also that S is the only ∇ -flat bilinear form which restricts to S^o for $x = 1$. Last, notice that the coefficient of θ^n in $S(\varepsilon, \eta)$, $\varepsilon, \eta \in G_0$, depends only on the classes of ε and η in $G_0/\theta G_0$. We will denote it by $g([\varepsilon], [\eta])$. This defines a nondegenerate bilinear form on $G_0/\theta G_0$ (see [11, p. 211]).

5 Application to the construction of Frobenius type structures and Frobenius manifolds

5.1 Frobenius type structures

In our situation, a Frobenius type structure on X is a tuple (see also [5] and [8])

$$(X, E, \nabla, R_0, R_\infty, \Phi, g)$$

where

- E is a $\mathbb{C}[x, x^{-1}]$ -free module,
- R_0 and R_∞ are $\mathbb{C}[x, x^{-1}]$ -linear endomorphisms of E ,
- $\Phi : E \rightarrow \Omega^1(X) \otimes E$ is a $\mathbb{C}[x, x^{-1}]$ -linear map,
- g is a metric on E , i.e. a $\mathbb{C}[x, x^{-1}]$ -bilinear form, symmetric and nondegenerate,
- ∇ is a connection on E ,

these objects satisfying the relations

$$\nabla^2 = 0, \nabla(R_\infty) = 0, \Phi \wedge \Phi = 0, [R_0, \Phi] = 0,$$

$$\nabla(\Phi) = 0, \nabla(R_0) + \Phi = [\Phi, R_\infty],$$

$$\nabla(g) = 0, \Phi^* = \Phi, R_0^* = R_0, R_\infty + R_\infty^* = rId$$

for a suitable constant r , $*$ denoting as above the adjoint with respect to g .

Keep the notations of section 3.1. The basis ω gives an extension of G_0 as a trivial bundle $\mathcal{G}$ on $\mathbb{P}^1 \times X$ (the module of its global sections is generated by $\omega_0, \dots, \omega_{\mu-1}$) equipped with a connection with logarithmic pole at $\tau := \theta^{-1} = 0$ and pole of Poincaré rank less or equal to one at $\theta = 0$. Define $E := i_{\{\theta=0\}} \mathcal{G}$, $E_\infty := i_{\{\tau=0\}} \mathcal{G}$ (E and E_∞ are canonically isomorphic) and, for $i, j = 0, \dots, \mu - 1$, $[\]$ denoting the class in E ,

- $R_0[\omega_i] := [\theta^2 \nabla_{\partial_\theta} \omega_i]$,
- $g([\omega_i], [\omega_j]) := \theta^{-n} S(\omega_i, \omega_j)$,
- $\Phi_\xi[\omega_i] := [\theta \nabla_{\partial_\xi} \omega_i]$ for any vector field ξ on X .

The connection ∇ and the endomorphism R_∞ are defined analogously, using the restriction E_∞ , $[\]$ denoting now the class in E_∞ ,

- $R_\infty[\omega_i] := [-\nabla_{\tau \partial_\tau} \omega_i]$,
- $\nabla_\xi[\omega_i] := [\nabla_{\partial_\xi} a]$.

Corollary 5.1.1 *The tuple $(X, E, R_0, R_\infty, \Phi, \nabla, g)$ is a Frobenius type structure on $X := \mathbb{C}^*$.*

Proof. This follows from proposition 3.1.3 and lemma 4.1.1 (see [11, Chapitre V, 2]). $\square$

5.2 Frobenius manifolds "in family"

Recall that Δ denotes the open disc in $\mathbb{C}$ of radius 1, centered at $x = 1$. Corollary 5.1.1 gives also an analytic Frobenius type structure

$$\mathcal{F} = (\Delta, E^{an}, R_0^{an}, R_\infty, \Phi^{an}, \nabla^{an}, g^{an})$$

on the *simply connected domain* Δ . Let ω_0^{an} be the class of ω_0 in E^{an} : ω_0^{an} is ∇^{an} -flat because $R(\omega_0) = 0$ (see remark 3.2.6). The universal deformations and the period maps that we will consider are the ones defined in [5] and [8].

Lemma 5.2.1 (1) *The Frobenius type structure $\mathcal{F}$ has a universal deformation*

$$\tilde{\mathcal{F}} = (N, \tilde{E}^{an}, \tilde{R}_0^{an}, \tilde{R}_\infty, \tilde{\Phi}^{an}, \tilde{\nabla}^{an}, \tilde{g}^{an})$$

parametrized by $N := \Delta \times (\mathbb{C}^{\mu-1}, 0)$.

(2) *The period map defined by the $\tilde{\nabla}^{an}$ -flat extension of ω_0^{an} to $\tilde{\mathcal{F}}$ is an isomorphism which makes N a Frobenius manifold.*

Proof. (1) We can use the adaptation of [8, Theorem 2.5] given in [5, Section 6] because $\omega_0^{an}, R_0^{an}(\omega_0^{an}), \dots, (R_0^{an})^{\mu-1}(\omega_0^{an})$ generate E^{an} and because $u_0 := 1/u_1^{w_1} \cdots u_n^{w_n}$ is not equal to zero in E^{an} . (2) follows from (1) (see e.g. [8, Theorem 4.5]). $\square$

The previous construction can be also done in the same way "point by point" (see [7] and [8] and the references therein) and, as quoted in the introduction, this is the classical point of view. Indeed, let $x \in \Delta$ and put $F_x := F(\cdot, x)$. One can attach to the Laurent polynomial F_x a Frobenius type structure on a point (see section 6.2 below) $\mathcal{F}_x^{pt}$, a universal deformation $\tilde{\mathcal{F}}_x^{pt}$ of it and finally, with the help of the section ω_0 , a Frobenius structure on $M := (\mathbb{C}^\mu, 0)$. We will call it "the Frobenius structure attached to F_x ". Let $\mathcal{F}_x$ (*resp.* $\tilde{\mathcal{F}}_x$) be the germ of $\mathcal{F}$ (*resp.* $\tilde{\mathcal{F}}$) at $x \in \Delta$ (*resp.* $(x, 0)$).

Proposition 5.2.2 (1) *The deformations $\tilde{\mathcal{F}}_x$ and $\tilde{\mathcal{F}}_x^{pt}$ are isomorphic.*

(2) *The period map defined by the flat extension of ω_0^{an} to $\tilde{\mathcal{F}}_x$ is an isomorphism. This yields a Frobenius structure on M which is isomorphic to the one attached to F_x .*

Proof. Note first that $\tilde{\mathcal{F}}_x^{pt}$ is a deformation of $\mathcal{F}_x$: this follows from the fact that u_0 does not belong to the Jacobian ideal of f : see [5, section 7]. Better, $\tilde{\mathcal{F}}_x^{pt}$ is a universal deformation of $\mathcal{F}_x$ because $\mathcal{F}_x$ is a deformation of $\mathcal{F}_x^{pt}$. This gives (1) because, by definition, two universal deformations of a same Frobenius type structure are isomorphic. (2) is then clear. $\square$

As a consequence, the universal deformations $\tilde{\mathcal{F}}_x^{pt}$, $x \in \Delta$, are the germs of a same section, namely $\tilde{\mathcal{F}}$. Thus, the Frobenius structure attached to F_{x_1} , $x_1 \in \Delta$, can be seen as an analytic continuation of the one attached to F_{x_0} , $x_0 \in \Delta$.

6 Limits

Our goal is now to define a canonical limit, as x approaches 0, of the Frobenius type structure constructed in corollary 5.1.1. We will use Deligne's canonical extensions.

6.1 Résumé: the canonical extensions at $x = 0$

Recall the lattices defined $\mathcal{L}^\varphi$ and $\mathcal{L}^\psi$ defined in remark 3.2.6. Put

$$\overline{\mathcal{L}}^\varphi := \mathcal{L}^\varphi / x\mathcal{L}^\varphi.$$

This is a free $\mathbb{C}[\theta, \theta^{-1}]$ -module of rank μ , equipped with a connection $\overline{\nabla}_{\partial_\theta}$ (induced by ∇_{∂_θ}). In the sequel, $\overline{\omega}^\varphi$ will denote the basis of $\overline{\mathcal{L}}^\varphi$ induced by ω^φ . Recall also that $\overline{A}_0^\varphi$ denotes the value of $A_0^\varphi(x)$ at $x = 0$: it is a $\mu \times \mu$ Jordan matrix. The following theorem summarizes the results obtained in the previous sections:

Theorem 6.1.1 1) $\mathcal{L}^\varphi$ is equipped with a connection

$$\nabla : \mathcal{L}^\varphi \rightarrow \theta^{-1}\Omega_{\mathbb{C}^* \times \mathbb{C}}(\log(\mathbb{C}^* \times \{0\})) \otimes \mathcal{L}^\varphi.$$

The matrix of $x\nabla_{\partial_x}$ in the basis ω^φ takes the form

$$-\frac{A_0^\varphi(x)}{\mu\theta} + \text{diag}(0, \dots, 0, \frac{s_{n+1}}{\mu}, \dots, \frac{s_{\mu-1}}{\mu})$$

and the one of ∇_{∂_θ} takes the form

$$\frac{A_0^\varphi(x)}{\theta^2} + \frac{A_\infty}{\theta}.$$

2) $x\nabla_{\partial_x}$ induces a map on $\overline{\mathcal{L}}^\varphi$ whose matrix, in the basis $\overline{\omega}^\varphi$, takes the form

$$-\frac{\overline{A}_0^\varphi}{\mu\theta} + \text{diag}(0, \dots, 0, \frac{s_{n+1}}{\mu}, \dots, \frac{s_{\mu-1}}{\mu}).$$

Its eigenvalues are contained in $[0, 1[$.

3) The matrix of $\overline{\nabla}_{\partial_\theta}$, acting on $\overline{\mathcal{L}}^\varphi$, takes the form, in the basis $\overline{\omega}^\varphi$,

$$\frac{\overline{A}_0^\varphi}{\theta^2} + \frac{A_\infty}{\theta}.$$

□

Corollary 6.1.2 $\mathcal{L}^\varphi$ is Deligne's canonical extension of the bundle G to $\mathbb{C}^* \times \mathbb{C}$ such that the eigenvalues of the residue of ∇_{∂_x} are contained in $[0, 1[$.

Analogous statements for $\overline{\mathcal{L}}^\psi := \mathcal{L}^\psi / x\mathcal{L}^\psi$ (replace $[0, 1[$ by $]-1, 0]$). $\overline{\mathcal{L}}^\varphi$ is the space of the "vanishing cycles" and $\overline{\mathcal{L}}^\psi$ is the one of the "nearby cycles". More generally, and after a base change of matrix $x^r I$, $r \in \mathbb{Z}$, the lattice $\mathcal{L}^\varphi$ (resp. $\mathcal{L}^\psi$) gives the canonical extensions whose eigenvalues are contained in $[k, k+1[$ (resp. $]k, k+1]$).

6.2 Limits of Frobenius type structures

Ideally, the limit of our Frobenius type structure as x approaches 0 should be a *Frobenius type structure on a point* that is a tuple

$$(E^{lim}, R_0^{lim}, R_\infty^{lim}, g^{lim})$$

where E^{lim} is a finite dimensional vector space over $\mathbb{C}$, g^{lim} is a symmetric and nondegenerate bilinear form on E^{lim} , R_0^{lim} and R_∞^{lim} being two endomorphisms of E^{lim} satisfying $(R_0^{lim})^* = R_0^{lim}$ and $R_\infty^{lim} + (R_\infty^{lim})^* = rId$ for a suitable complex number r , $*$ denoting the adjoint with respect to g . It turns out that our limit will be defined with the help of the graded module $gr^V(\overline{\mathcal{L}}^\varphi)$ associated with the Kashiwara-Malgrange V -filtration at $x = 0$.

6.2.1 The V -filtration at $x = 0$

Recall the basis $\omega^\varphi = (\omega_0^\varphi, \dots, \omega_{\mu-1}^\varphi)$ of G_0 over $\mathbb{C}[\theta, x, x^{-1}]$ (it is thus also a basis of G over $\mathbb{C}[\theta, \theta^{-1}, x, x^{-1}]$) defined in section 3.2.1. Put $v(\omega_0^\varphi) = \dots = v(\omega_n^\varphi) = 0$ and, for $k = n+1, \dots, \mu-1$, $v(\omega_k^\varphi) = s_k/\mu$. Define, for $0 \leq \alpha < 1$,

$$V^\alpha G = \sum_{\alpha \leq v(\omega_k^\varphi)} \mathbb{C}\{x\}[\theta, \theta^{-1}]\omega_k^\varphi + x \sum_{\alpha > v(\omega_k^\varphi)} \mathbb{C}\{x\}[\theta, \theta^{-1}]\omega_k^\varphi,$$

$$V^{>\alpha} G = \sum_{\alpha < v(\omega_k^\varphi)} \mathbb{C}\{x\}[\theta, \theta^{-1}]\omega_k^\varphi + x \sum_{\alpha \geq v(\omega_k^\varphi)} \mathbb{C}\{x\}[\theta, \theta^{-1}]\omega_k^\varphi$$

and $V^{\alpha+p}G = x^p V^\alpha G$ for $p \in \mathbb{Z}$ and $\alpha \in [0, 1[$. This gives a *decreasing* filtration $V^\bullet$ of G by $\mathbb{C}\{x\}[\theta, \theta^{-1}]$ -submodules such that

$$V^\alpha G = \mathbb{C}[\theta, \theta^{-1}] < \overline{\omega}_k^\varphi | v(\omega_k^\varphi) = \alpha > + V^{>\alpha} G.$$

Notice that $\mathcal{L}^\varphi = V^0 G$ and that $\overline{\mathcal{L}}^\varphi = V^0 G / V^1 G$. We will put $H^\alpha := V^\alpha G / V^{>\alpha} G$ and $H = \bigoplus_{\alpha \in [0, 1[} H^\alpha$.

Lemma 6.2.1 (1) For each α , $(x\nabla_{\partial_x} - \alpha)$ is nilpotent on H^α .

(2) Let N be the nilpotent endomorphism of H which restricts to $(x\nabla_{\partial_x} - \alpha)$ on H^α . Its Jordan blocks are in one to one correspondance with the maximal constant sequences in $\mathcal{S}_w$ and the corresponding sizes are the same.

(3) Let e_k be the class of ω_k^φ in H . Then $e = (e_0, \dots, e_{\mu-1})$ is a basis of H over $\mathbb{C}[\theta, \theta^{-1}]$.

Proof. (1) It suffices to prove the assertion for $\alpha \in [0, 1[$. By theorem 6.1.1, we have

$$x\nabla_{\partial_x} \omega_k^\varphi = -\frac{1}{\theta} \omega_{k+1}^\varphi$$

for $k = 0, \dots, n-1$ and $x\nabla_{\partial_x} \omega_n^\varphi \in V^{>0} G$. Moreover, for $k = n+1, \dots, \mu-2$ we have

$$(x\nabla_{\partial_x} - \frac{s_k}{\mu}) \omega_k^\varphi = -\frac{1}{\theta} \omega_{k+1}^\varphi$$

and this is equal to 0 in $H^{v(\omega_k^\varphi)}$ if $s_{k+1} > s_k$. Last,

$$(x\nabla_{\partial_x} - \frac{s_{\mu-1}}{\mu})\omega_{\mu-1}^\varphi \in x \sum_{v(\omega_{\mu-1}^\varphi) \geq v(\omega_k^\varphi)} \mathbb{C}\{x\}\bar{\omega}_k^\varphi \in V^{>s_{\mu-1}/\mu}G.$$

(2) follows from (1) and (3) follows from the definition of $V^\bullet$. $\square$

Remark 6.2.2 Let B be the matrix of N in the basis e : we have $B_{i,j} = 0$ if $i \neq j+1$, $B_{i+1,i} = -\frac{1}{\theta}$ if $\alpha_i = \alpha_{i-1} + 1$ (equivalently if $s_i = s_{i-1}$) and $B_{i+1,i} = 0$ if $\alpha_i \neq \alpha_{i-1} + 1$.

Example 6.2.3 (1) $n = 2$ and $w_1 = w_2 = 2$: one has $\mathcal{S}_w = (0, 0, 0, \frac{5}{2}, \frac{5}{2})$ so that N has one Jordan block of size 3 and one Jordan block of size 2. Its matrix in the basis e is

$$-\frac{1}{\theta} \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix}.$$

(2) $w_1 = \dots = w_n = 1$: one has $\mathcal{S}_w = (0, \dots, 0)$ so that N has one Jordan block of size $\mu = n + 1$. Its matrix in the basis e is

$$-\frac{1}{\theta} \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \dots & 1 & 0 \end{pmatrix}.$$

Corollary 6.2.4 The filtration $V^\bullet$ is the Kashiwara-Malgrange filtration at $x = 0$.

Proof. By lemma 6.2.1, our filtration satisfies all the characterizing properties of the Kashiwara-Malgrange filtration at $x = 0$. $\square$

6.2.2 The canonical limit Frobenius type structure

The module H is free over $\mathbb{C}[\theta, \theta^{-1}]$ and is equipped with a connection ∇ whose matrix in the basis e takes the form

$$(\frac{[A_0]}{\theta} + [A_\infty])\frac{d\theta}{\theta}$$

where $[A_0] = -\mu\theta B$ (as above, B is the matrix of N in the basis e) and where $[A_\infty]$ is the diagonal matrix with eigenvalues $(\alpha_0, \dots, \alpha_{\mu-1})$. Let H_0 be the $\mathbb{C}[\theta]$ -submodule of H generated by e and define

$$k : H_0 \times H_0 \rightarrow \mathbb{C}[\theta]\theta^n$$

by

$$k(e_k, e_{n-k}) = \theta^n$$

for $k = 0, \dots, n$,

$$k(e_k, e_{\mu+n-k}) = \theta^n$$

for $k = n+1, \dots, \mu-1$ and $k(e_i, e_j) = 0$ otherwise. Last, put $E^{lim} = H_0/\theta H_0$. E^{lim} is thus equipped with two endomorphisms R_0^{lim} and R_∞^{lim} and with a non-degenerate bilinear form g^{lim} : if $\bar{e}_i$ denotes the class of e_i in E^{lim} , R_0^{lim} (resp. R_∞^{lim}) is the endomorphism of E^{lim} whose matrix is $[A_0]$ (resp. $[A_\infty]$) in the basis $\bar{e}$ and g^{lim} is obtained from k as in the end of section 4.

Theorem 6.2.5 *The tuple*

$$(E^{lim}, R_0^{lim}, R_\infty^{lim}, g^{lim})$$

is a Frobenius type structure on a point.

Proof. It is enough to show that $(R_0^{lim})^* = R_0^{lim}$ and that $R_\infty^{lim} + (R_\infty^{lim})^* = nId$. To make the proof readable, we write e_i instead of $\bar{e}_i$, R_0 instead of R_0^{lim} etc... We will repeatedly use lemma 2.1.1, corollary 2.1.2 and lemma 6.2.1.

(a) Let us show first that $R_\infty + (R_\infty)^* = nId$:

(i) assume $0 \leq i \leq n$: we have $g(R_\infty(e_i), e_j) = \alpha_i g(e_i, e_j)$ so that

$$g(R_\infty(e_i), e_j) = \alpha_i$$

if $i+j = n$ and $g(R_\infty(e_i), e_j) = 0$ otherwise. In the same way,

$$g(e_i, R_\infty(e_j)) = \alpha_j$$

if $i+j = n$ and $g(e_i, R_\infty(e_j)) = 0$ otherwise. If $i+j = n$, one has $\alpha_j = \alpha_{n-i} = n-i - s_{n-i} = n-i = n - \alpha_i$.

(ii) Assume now $n+1 \leq i \leq \mu-1$: we have

$$g(R_\infty(e_i), e_j) = \alpha_i$$

if $i+j = \mu+n$ and $g(R_\infty(e_i), e_j) = 0$ otherwise. In the same way,

$$g(e_i, R_\infty(e_j)) = \alpha_j$$

if $i+j = \mu+n$ and $g(e_i, R_\infty(e_j)) = 0$ otherwise. If $i+j = \mu+n$, one has

$$\alpha_j = \alpha_{\mu+n-i} = \mu+n-i - s_{\mu+n-i} = \mu+n-i + (s_i - \mu) = n - \alpha_i.$$

(b) Let us show now that $(R_0)^* = R_0$.

(i) Assume that $0 \leq i \leq n-1$. If $i+j+1 = n$ (thus $0 \leq j \leq n-1$), one has

$$g(R_0(e_i), e_j) = g(e_{i+1}, e_j) = 1$$

and

$$g(e_i, R_0(e_j)) = g(e_i, e_{j+1}) = 1.$$

If $i+j+1 \neq n$, one always has

$$g(R_0(e_i), e_j) = g(e_i, R_0(e_j)) = 0.$$

(ii) Assume that $i = n$. Then $g(R_0(e_n), e_j) = 0$ because $R_0(e_n) = 0$ by lemma 6.2.1 and $g(e_n, R_0(e_j)) = 0$ because e_0 does not belong to the image of R_0 .

(iii) Assume $n + 1 \leq i \leq \mu - 1$.

If $i + 1 + j = \mu + n$ then $s_{i+1} = s_i$ if and only if $s_{j+1} = s_j$ (because $s_{\mu+n-i} = \mu - s_i$). If it is the case, one has

$$g(R_0(e_i), e_j) = g(e_{i+1}, e_j) = 1$$

and

$$g(e_i, R_0(e_j)) = g(e_i, e_{j+1}) = 1.$$

If $i + 1 + j = \mu + n$ but $s_{i+1} \neq s_i$ (and thus $s_{j+1} \neq s_j$), one has $R_0(e_i) = 0$ and $R_0(e_j) = 0$ so that

$$g(R_0(e_i), e_j) = g(e_i, R_0(e_j)) = 0.$$

If $i + 1 + j \neq \mu + n$, one always has

$$g(R_0(e_i), e_j) = g(e_i, R_0(e_j)) = 0.$$

□

Remark 6.2.6 *The conclusion of the previous theorem is not always true if we do not consider the graded module $H := gr^V(\mathcal{L}^\varphi/x\mathcal{L}^\varphi)$. Indeed, if we work directly on $\overline{\mathcal{L}}^\varphi := \mathcal{L}^\varphi/x\mathcal{L}^\varphi$ we can define, in the same way as above, the tuple*

$$(\mathcal{E}, \mathcal{R}_0, \mathcal{R}_\infty, \mathcal{G})$$

where

- $\mathcal{E} = \overline{\mathcal{L}}^\varphi/\theta\overline{\mathcal{L}}^\varphi$,
- $\mathcal{G}$ is the symmetric and nondegenerate bilinear form on $\mathcal{E}$ defined by $\mathcal{G}(e'_k, e'_{n-k}) = 1$ for $k = 0, \dots, n$, $\mathcal{G}(e'_k, e'_{\mu+n-k}) = 1$ for $k = n+1, \dots, \mu-1$ and $\mathcal{G}(e'_i, e'_j) = 0$ otherwise, e'_k denoting the class of ω_k^φ in $\mathcal{E}$,
- $\mathcal{R}_0$ (resp. $\mathcal{R}_\infty$) is the endomorphism of $\mathcal{E}$ whose matrix is $\overline{A}_0^\varphi$ (resp. A_∞) in the basis $e' = (e'_0, \dots, e'_{\mu-1})$.

The point is that this tuple is a Frobenius type structure on a point if and only if $\mu = n + 1$: for instance, if $\mu \geq n + 2$, we have $\mathcal{G}(\mathcal{R}_0(e'_n), e'_{\mu-1}) = 1$ but $\mathcal{G}(e'_n, \mathcal{R}_0(e'_{\mu-1})) = 0$ so that $(\mathcal{R}_0)^* \neq \mathcal{R}_0$. This symmetry default shows that the tuple $(\mathcal{E}, \mathcal{R}_0, \mathcal{R}_\infty, \mathcal{G})$ is not a Frobenius type structure. The case $\mu = n + 1$ is directly checked.

6.3 Pre-primitive sections "at the limit"

We will say that an element e of a μ -dimensional vector space E over $\mathbb{C}$, equipped with two endomorphisms A and B , is a *pre-primitive section* of the triple (E, A, B) if $(e, A(e), \dots, A^{\mu-1}(e))$ is a basis of E over $\mathbb{C}$ and that e is *homogeneous* if it is an eigenvector of B . Let

$$(E^{lim}, R_0^{lim}, R_\infty^{lim}, g^{lim})$$

be the limit Frobenius structure given by theorem 6.2.5. Recall that $\overline{e}_0$ denotes the class of ω_0^φ in E^{lim} .

Lemma 6.3.1 (1) $\bar{e}_0$ is a homogeneous section of the triple $(E^{lim}, R_0^{lim}, R_\infty^{lim})$,
(2) $\bar{e}_0$ is a pre-primitive section of the triple $(E^{lim}, R_0^{lim}, R_\infty^{lim})$ if and only if $\mu = n+1$. If $\mu \geq n+2$, this triple has no pre-primitive section at all.

Proof. Obvious, except the last assertion: this follows from the fact that if $\mu \geq n+2$, R_0^{lim} has at least two Jordan blocks for the same eigenvalue 0 (see lemma 6.2.1). $\square$

Corollary 6.3.2 $\bar{e}_0$ is a pre-primitive and homogeneous section of the limit Frobenius type structure $(E^{lim}, R_0^{lim}, R_\infty^{lim}, g^{lim})$ if and only if $\mu = n+1$.

6.4 A canonical limit Frobenius manifold

What do we need to construct a Frobenius manifold? In general, a Frobenius type structure and a pre-primitive and homogeneous section of it: the main point is that these two objects give a unique (up to isomorphism) Frobenius manifold, see for instance [8] and the references to B. Dubrovin and B. Malgrange therein.

We assume here that $\mu = n+1$: theorem 6.2.5 gives a canonical limit Frobenius type structure and corollary 6.3.2 a pre-primitive and homogeneous section of it, so, as explained above, we get in this case a canonical (limit) Frobenius manifold. We can give in this case a precise description of it: recall that $[A_0]$

$$[A_0] = (n+1) \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \vdots & \cdots & 1 & 0 \end{pmatrix}, \quad A_\infty = \text{diag}(0, 1, \dots, n).$$

Let $x = (x_1, \dots, x_\mu)$ be a system of coordinates on $M = (\mathbb{C}^\mu, 0)$.

Lemma 6.4.1 *There exists a unique tuple of matrices*

$$(\tilde{A}_0(x), A_\infty, \tilde{C}_1(x), \dots, \tilde{C}_\mu(x))$$

such that $\tilde{A}_0(0) = [A_0]$, $(\tilde{C}_i)_{i1} = -1$ for all $i = 1, \dots, \mu$ and satisfying the relations

$$\begin{aligned} \frac{\partial \tilde{C}_i}{\partial x_j} &= \frac{\partial \tilde{C}_j}{\partial x_i} \\ [\tilde{C}_i, \tilde{C}_j] &= 0 \\ [\tilde{A}_0(x), \tilde{C}_i] &= 0 \\ \frac{\partial \tilde{A}_0}{\partial x_i} + \tilde{C}_i &= [A_\infty, \tilde{C}_i] \end{aligned}$$

for all $i, j = 1, \dots, \mu$. Precisely,

- (a) $\tilde{C}_1 = -I$,
- (b) $\tilde{C}_2 = -J$ where J denotes the nilpotent Jordan matrix of order μ ,
- (c) $\tilde{C}_i = -J^{i-1}$ for all $i = 1, \dots, \mu$.
- (d) $\tilde{A}_0(x) = -x_1 \tilde{C}_1 - \mu \tilde{C}_2 + x_3 \tilde{C}_3 + 2x_4 \tilde{C}_4 + \dots + (\mu-2)x_\mu \tilde{C}_\mu$.

Proof. It is clear that the given matrices satisfy the required relations. To show unicity, note that the matrices $\tilde{C}_i$ are determined by their first column because the matrix $\tilde{A}_0(0)$ is regular, $\bar{e}_0$ is pre-primitive and the matrices $\tilde{C}_i$ commute with $\tilde{A}_0(x)$. $\square$

This lemma means the following: the connection $\tilde{\nabla}$ on the bundle $\tilde{E} = \mathcal{O}_M \otimes E^{lim}$ whose matrix is

$$\left(\frac{\tilde{A}_0(x)}{\theta} + A_\infty\right)\frac{d\theta}{\theta} + \frac{\sum_{i=1}^\mu \tilde{C}_i dx_i}{\theta}$$

in the basis $\tilde{e} = (\tilde{e}_0, \dots, \tilde{e}_{\mu-1}) = (1 \otimes \bar{e}_0, \dots, 1 \otimes \bar{e}_{\mu-1})$ of $\tilde{E}$ is *flat*. The matrices $\tilde{C}_i$ are the matrices of the covariant derivatives $\tilde{\nabla}_{\partial_{x_i}}$. $\tilde{\nabla}$ will denote the connection on $\tilde{E}$ whose matrix is zero in the basis $\tilde{e}$: $\tilde{e}$ is thus the $\tilde{\nabla}$ -flat extension of $\bar{e}$. We get in this way a Frobenius type structure on M ,

$$(M, \tilde{E}, \tilde{\nabla}, \tilde{R}_0, \tilde{R}_\infty, \tilde{\Phi}, \tilde{g}).$$

Corollary 6.4.2 *Assume that $\mu = n + 1$.*

(1) *The period map*

$$\varphi_{\tilde{e}_0} : TM \rightarrow \tilde{E},$$

$\varphi_{\tilde{e}_0}(\xi) = -\tilde{\Phi}_\xi(\tilde{e}_0)$, *is an isomorphism and $\tilde{e}_0$ is a homogeneous section of $\tilde{E}$, that is an eigenvector of $\tilde{R}_\infty$.*

(2) *The section $\tilde{e}_0$ defines, through the period map $\varphi_{\tilde{e}_0}$ a Frobenius structure on M which makes M a Frobenius manifold for which:*

- (a) *the coordinates $(x_1, \dots, x_\mu)$ are ∇ -flat: one has $\nabla \partial_{x_i} = 0$ for all $i = 1, \dots, \mu$,*
- (b) *the product is constant in flat coordinates: $\partial_{x_i} * \partial_{x_j} = \partial_{x_{i+j-1}}$ if $i + j - 1 \leq \mu$, 0 otherwise,*
- (c) *the potential Ψ is a polynomial of degree less or equal to 3: $\Psi = \sum_{i,j, i+j \leq \mu+1} a_{ij} x_i x_j x_{\mu+2-i-j}$, up to a polynomial of degree less or equal to 2,*
- (d) *the Euler vector field is $E = x_1 \partial_{x_1} + (n+1) \partial_{x_2} - x_3 \partial_{x_3} - \dots - (n-1) x_{n+1} \partial_{x_{n+1}}$,*
- (e) *the potential Ψ is, up to polynomials of degree less or equal to 2, Euler-homogeneous of degree $4 - \mu$:*

$$E(\Psi) = (4 - \mu)\Psi + G(x_1, \dots, x_\mu)$$

where G is a polynomial of degree less or equal to 2.

Proof. (1) Follows from the choice of the first columns of the matrices $\tilde{C}_i$: indeed, the period map $\varphi_{\tilde{e}_0}$ is defined by $\varphi_{\tilde{e}_0}(\partial_{x_i}) = -\tilde{C}_i(\tilde{e}_0) = \tilde{e}_{i-1}$, and it is of course an isomorphism. Last, $\tilde{e}_0$ is homogeneous because $\bar{e}_0$ is so. Let us show (2): the isomorphism $\varphi_{\tilde{e}_0}$ brings on TM the structures on $\tilde{E}$: (a) follows from the fact that the first column of the matrices $\tilde{C}_i$ are constant and (b) from the fact that the matrices $\tilde{C}_i$ are constant because, by the definition of the product, $\varphi_{\tilde{e}_0}(\partial_{x_i} * \partial_{x_j}) = \tilde{C}_i(\tilde{C}_j(\tilde{e}_0))$; (c) follows from (b) because, in flat coordinates,

$$g(\partial_{x_i} * \partial_{x_j}, \partial_{x_k}) = \frac{\partial^3 \Psi}{\partial x_i \partial x_j \partial x_k}.$$

Last, (d) follows from the definition of $\tilde{A}_0(x)$ and (e) is a consequence of (c) and (d). $\square$

Of course, the period map can be an isomorphism for other choices of the first columns of the matrices C_i : whatever happens, the resulting Frobenius manifolds will be isomorphic to the one given by the corollary. Indeed, the Frobenius type structure

$$(M, \tilde{E}, \tilde{\nabla}, \tilde{R}_0, \tilde{R}_\infty, \tilde{\Phi}, \tilde{g})$$

is a *universal* deformation of the limit Frobenius type structure $(E^{lim}, R_0^{lim}, R_\infty^{lim}, g^{lim})$ given by theorem 6.2.5 (see [8]). We will thus call the Frobenius manifold given by the corollary the *canonical limit Frobenius manifold*.

Remark 6.4.3 *If $\mu \geq n+2$, that is if there exists an w_i such that $w_i \geq 2$, we still have a canonical limit Frobenius type structure, but no pre-primitive section of it so that the results in [8] do not apply. In particular, we do not know if one can find matrices as lemma 6.4.1 (this problem is not obvious, even for the simplest examples, see for instance example 2.2.1 (1)), that is if the limit Frobenius type structure and the form $\bar{e}_0$ (or any other) give as above a (limit) Frobenius manifold through the period map. Even if it happens to be the case, the previous construction gives then a lot of (limit) Frobenius manifolds and there is no way to compare them (we do not have any kind of unicity here).*

7 Logarithmic Frobenius type structures and logarithmic Frobenius manifolds: an example and some remarks

Proposition 3.1.3 suggests that we are not so far from a logarithmic Frobenius type structure in the sense of [10, Definition 1.6] and one could expect at the end a logarithmic Frobenius manifold, see [10, Definition 1.4]. Some remarks are in order.

7.1 Logarithmic Frobenius type structures

A Frobenius type structure with *logarithmic pole along $\{x = 0\}$* (for short, a logarithmic Frobenius type structure) is a tuple

$$(E^{log}, \{0\}, \nabla, R_0, R_\infty, \Phi, g)$$

where

- E^{log} is a $\mathbb{C}[x]$ -free module,
- R_0 and R_∞ are $\mathbb{C}[x]$ -linear endomorphisms of E^{log} ,
- $\Phi : E^{log} \rightarrow \Omega^1(\log(\{x = 0\})) \otimes E^{log}$ is a $\mathbb{C}[x]$ -linear map,
- g is a metric on E^{log} , i.e a $\mathbb{C}[x]$ -bilinear form, symmetric and non-degenerate,
- ∇ is a connection on E^{log} with logarithmic pole along $\{x = 0\}$,

these object satisfying the compatibility relations of section 5.1. One can also define in an obvious way a logarithmic Frobenius type structure *without metric*.

The main point is to construct E^{log} : in our situation, it will be obtained from an extension of G_0 as a free $\mathbb{C}[x, \theta]$ -module (and not from a canonical extension of G as before). As pointed me out by C. Sevenheck, we can use for instance the $\mathbb{C}[x, \theta]$ -submodule of G_0 generated by $\omega_0^\varphi, \dots, \omega_{\mu-1}^\varphi$ which is a lattice “in x ” of G_0 . We will denote it by $\mathcal{L}_0^\varphi$. Let $\mathcal{L}_\infty^\varphi$ be the $\mathbb{C}[x, \tau]$ -module generated by $\omega_0^\varphi, \dots, \omega_{\mu-1}^\varphi$ (as usual $\tau = \theta^{-1}$).

Lemma 7.1.1 $\mathcal{L}_0^\varphi$ is equipped with a connection

$$\nabla : \mathcal{L}_0^\varphi \rightarrow \theta^{-1} \Omega_{\mathbb{C} \times \Delta}(\log((\{0\} \times \mathbb{C}) \cup (\mathbb{C} \times \{0\}))) \otimes \mathcal{L}_0^\varphi.$$

The matrix of $x\nabla_{\partial_x}$ in the basis ω^φ of $\mathcal{L}_0^\varphi$ takes the form

$$-\frac{A_0^\varphi(x)}{\mu\theta} + \text{diag}(0, \dots, 0, \frac{s_{n+1}}{\mu}, \dots, \frac{s_{\mu-1}}{\mu})$$

and the one of ∇_{∂_θ} takes the form

$$\frac{A_0^\varphi(x)}{\theta^2} + \frac{A_\infty}{\theta}.$$

Proof. Follows from proposition 3.2.1. □

Define $E^{log} = \mathcal{L}_0^\varphi / \theta \mathcal{L}_0^\varphi$. One could imagine that the counterpart of corollary 5.1.1: indeed, define, for $i = 0, \dots, \mu - 1$,

- $R_0[\omega_i^\varphi] := [\theta^2 \nabla_{\partial_\theta} \omega_i^\varphi]$,
- $\Phi_\xi[\omega_i^\varphi] := [\theta \nabla_{\partial_\xi} \omega_i^\varphi]$ for any logarithmic vector field $\xi \in \text{Der}(\log\{x = 0\})$,

and, using the restriction of $\mathcal{L}_\infty^\varphi$ to $\tau = 0$,

- $R_\infty[\omega_i^\varphi] := [-\nabla_{\tau \partial_\tau} \omega_i^\varphi]$,
- $\nabla_\xi[\omega_i^\varphi] = [\nabla_{\partial_\xi} \omega_i^\varphi]$ for any logarithmic vector field $\xi \in \text{Der}(\log\{x = 0\})$.

In order to define the ‘metric’, recall that (see section 4)

$$S(\omega_0^\varphi, \omega_n^\varphi) = \theta^n,$$

$$S(\omega_k^\varphi, \omega_{n-k}^\varphi) = \theta^n$$

for $k = 1, \dots, n - 1$,

$$S(\omega_k^\varphi, \omega_{\mu+n-k}^\varphi) = x\theta^n$$

for $k = n + 1, \dots, \mu - 1$ and

$$S(\omega_i^\varphi, \omega_j^\varphi) = 0$$

otherwise. Extend S to $\mathcal{L}_0^\varphi$: as above, we get a flat bilinear symmetric form g on E^{log} ,

$$g([\omega_i^\varphi], [\omega_j^\varphi]) := \theta^{-n} S(\omega_i^\varphi, \omega_j^\varphi).$$

The main point is that, of course, g which is not non-degenerate, unless $\mu = n + 1$.

Corollary 7.1.2 (1) The tuple $(E^{\log}, \{0\}, R_0, R_\infty, \Phi, \nabla, g)$ is a logarithmic Frobenius type structure if $\mu = n + 1$.
(2) The tuple $(E^{\log}, \{0\}, R_0, R_\infty, \Phi, \nabla)$ is a logarithmic Frobenius type structure without metric if $\mu \geq n + 2$.

Proof. The previous lemma finally gives a $\log(\{x = 0\}) - trTLEP$ -structure (see [10, Definition 1.8]) if $\mu = n + 1$ and we use the 1-1 correspondance between such structures and logarithmic Frobenius type structures given by [10, Proposition 1.10]. $\square$

Remark 7.1.3 (1) We can also consider the lattice $\mathcal{L}_0$ (resp. $\mathcal{L}_0^\psi$), defined using the basis ω (resp. ω^ψ). We have

$$\mathcal{L}_0^\varphi \subset \mathcal{L}_0^\psi \subset \mathcal{L}_0.$$

Anyway, we will see below that, even if we forget the metric, the 'good' one to consider is $\mathcal{L}_0^\varphi$.

(2) In [4], and for $w_1 = \dots = w_n = 1$, another (and more intrinsic) extension of G_0 is considered, built with the help of logarithmic vector fields. I don't know for the moment how to compare it with $\mathcal{L}_0^\varphi$ and if one can define in this way extensions of G_0 if there exists an $w_i \geq 2$.

7.2 Construction of a logarithmic Frobenius manifold

A manifold M is a *Frobenius manifold with logarithmic poles along the divisor D* (for short a logarithmic Frobenius manifold) if $Der_M(\log D)$ is equipped with a metric, a multiplication and two (global) logarithmic vector fields (the unit e for the multiplication and the Euler vector field E), all these objects satisfying the usual compatibility relations (see [10, Definition 1.4]). We can also define a Frobenius manifold with logarithmic poles *without metric*: in this case, we still need a flat, torsionless connection, a symmetric Higgs field (that is a product) and two global logarithmic vector fields as before. Of course $D = \{x = 0\}$ in what follows. According to T. Reichelt [10, Theorem 1.12] the construction in section [8] can be adapted to get a logarithmic Frobenius manifold from a logarithmic Frobenius type structure: let $(E^{\log}, D, R_0, R_\infty, \Phi, \nabla, g)$ be a logarithmic Frobenius type structure, ω be a section of $E^{\log}$. Define

$$\varphi_\omega : Der_M(\log D) \rightarrow E^{\log},$$

by

$$\varphi_\omega(\xi) := -\Phi_\xi(\omega).$$

One says that ω satisfies

- (IC) if $\varphi_\omega|_0$ is injective,
- (GC) if $\omega|_0$ and its images under iteration of the maps $\Phi_\xi|_0$, $\xi \in Der_M(\log D)$, and $R_0|_0$ generate $E^{\log}|_0$,
- (EC) if ω is an eigenvector of R_∞ .

We will say that a section of $E^{\log}$ is *log-pre-primitive* (resp. *homogeneous*) if its restriction to $M - D$ is ∇ -flat and if it satisfies conditions (IC), (GC) (resp. (EC)). We now come back to the logarithmic Frobenius type structure given by corollary 7.1.2.

Lemma 7.2.1 *The (class of the) form ω_0^φ in $E^{\log}$ is log-pre-primitive and homogeneous.*

Proof. Follows from proposition 3.2.1: the flatness is given by the fact that $R^\varphi(\omega_0^\varphi) = 0$; conditions (IC) and (GC) hold because the matrix of $\Phi_{x\partial_x}$ is $-\frac{A_0^\varphi(x)}{\mu}$; last, we have $A_\infty(\omega_0^\varphi) = 0$ and this gives (EC). $\square$

Remark 7.2.2 *Assume that $\mu \geq n + 2$: the section ω_0 in $\mathcal{L}_0$ is flat but does not satisfy (IC) and the section ω_0^ψ in $\mathcal{L}_0^\psi$ is flat but does not satisfy (GC). In the former case, the only section which satisfies (IC), (EC) and (GC) is ω_1 but this one is not flat; in the latter case, the only section which satisfies (IC), (EC) and (GC) is ω_{n+1}^ψ but this one is not flat. This explains why we work with $\mathcal{L}_0^\varphi$.*

Corollary 7.2.3 *The log-pre-primitive and homogeneous section ω_0^φ together with the logarithmic Frobenius type structure $(E^{\log}, \{0\}, R_0, R_\infty, \Phi, \nabla, g)$ define a logarithmic Frobenius manifold if $\mu = n + 1$ and a logarithmic Frobenius manifold without metric if $\mu \geq n + 2$.*

Proof. Follows now from [10, theorem 1.12]. $\square$

If $\mu = n + 1$, one could expect an explicit description of the logarithmic Frobenius manifold obtained, as in section 6.4. Unfortunately, it is much more difficult and, except some trivial cases, I do not have results in this direction.

References

- [1] Barannikov, S.: Semi-infinite Hodge structures and mirror symmetry for projective spaces. Available at arXiv.math.AG/0010157.
- [2] Bolibruch, A.A : On isomonodromic confluences of Fuchsian systems. Proceedings of the Steklov Institute of Mathematics **221**, 1998, p. 117-132.
- [3] Coates, T., Corti, A., Lee, Y., Tseng, H.: The quantum orbifold cohomology of weighted projective space. Available at arXiv.math.AG/0608481.
- [4] de Gregorio, I., Mond, D., Sevenheck, C.: Linear free divisors and Frobenius manifolds. Available at arXiv:0802.4188v1.
- [5] Douai, A.: A canonical Frobenius structure. To appear in Math. Z. The publication is already available at www.springerlink.com (doi:10.1007/s00209-008-0344-3).
- [6] Douai, A., Sabbah, C.: Gauss-Manin systems, Brieskorn lattices and Frobenius structures I. Ann. Inst. Fourier **53-4**, 2003, p. 1055-1116.
- [7] Douai, A., Sabbah, C.: Gauss-Manin systems, Brieskorn lattices and Frobenius structures II. In: Frobenius Manifolds, C. Hertling and M. Marcolli (Eds.), Aspects of Mathematics E 36.
- [8] Hertling, C., Manin, Y.: Meromorphic connections and Frobenius manifolds. In: Frobenius Manifolds, C. Hertling and M. Marcolli (Eds.), Aspects of Mathematics E 36.

- [9] Mann, E.: Orbifold quantum cohomology of weighted projective spaces. Available at [arXiv.math.AG/0610965](https://arxiv.org/abs/math/0610965).
- [10] Reichelt, T.: A construction of Frobenius manifolds with logarithmic poles and applications. Available at [arXiv:0808.2698v1](https://arxiv.org/abs/0808.2698v1).
- [11] Sabbah, C.: Déformations isomonodromiques et variétés de Frobenius. Savoirs Actuels, CNRS Editions, Paris, 2002.